

Focused Critical Care Ultrasound Study (FOCCUS)

André Y. Denault, MD, PhD, ABIM-CCM, FASE

Cardiac Anesthesiologist, Montreal Heart Institute and Université de Montréal,
Critical Care Physician, Centre Hospitalier de l'Université de Montréal (CHUM)

Correspondence: Dr André Y. Denault, Department of Anesthesia, Montreal Heart Institute,
5000 Belanger Street, Montreal, Quebec H1T 1C8. Tel.: 514-376-3330 x3732; Fax: 514-
376-8784; email: andre.denault@umontreal.ca

Over the last decade, there have been significant developments in bedside echocardiography in the intensive care unit (ICU), the operating room (OR), the non-cardiac setting or the recovery room for hemodynamically unstable patients and patients with contraindications to transesophageal echocardiography (TEE). An approach combining both physical examination and bedside echography has improved the clinical diagnosis and management of acutely ill patients. Such point-of-care echographic examinations are usually "directed" or "focused" toward a specific clinical question and are significantly shorter in duration (less than six minutes in some studies) than conventional echocardiography.^{1,2} The goal of such an exam is not to perform a complete, comprehensive study, but rather to further enhance and extend the physical examination.

Transthoracic echocardiography (TTE) is non-invasive and more readily available than TEE and should, therefore, be the modality of choice to perform a goal-directed echocardiographic or echographic examination.³ So far, it has not been widely used in the ICU and the OR because only a few intensivists and anesthesiologists are TTE-trained.⁴ Nonetheless, over the last few years, bedside echocardiography has been increasingly performed by non-cardiologists, intensivists and anesthesiologists to provide diagnostic information not assessable by physical examination alone.⁵⁻⁷³ The TTE exam is called focused, goal-directed TTE, called FOCUS (FOcused Cardiac Ultrasound Study) or FOCCUS (FOcused Critical Care Ultrasound Study), is defined as a TTE performed with specific, limited objectives.¹⁰ The FOCUS objectives include the assessment of left ventricular (LV) function, right ventricular (RV) function, the assessment of the pericardial space for effusion and tamponade, and the assessment of volume status. Three views are used to evaluate cardiac function (Figure 1).




Figure 1 Basic FOCUS echocardiographic views. 1. Parasternal long-axis (A,B) and parasternal short-axis (C,D) views are shown. 2. Apical four-chamber view (E,F) and (3) Subcostal view (G,H) are shown. Abbreviations: Ao, aorta; LA, left atrium; FOCUS, focused cardiac ultrasound study; LV, left ventricle; RA, right atrium; RV, right ventricle. Source: Adapted from the FOCUS pocket guide, with permission from ICCU Imaging Inc. (From Denault A, Couture P, Vegas A, Buithieu J, Tardif JC. TEE multimedia manual 2nd edition.)

Additional FOCCUS objectives include the assessment of lung pathology⁸, abdominal fluid, vessels assessment, etc. Competency will eventually have to be established following predetermined levels of training (e.g., levels 1 to 4). Some training guidelines have already been proposed,⁹⁻¹¹ including a consensus between the American College of Chest Physicians and the Société de Réanimation de Langue Française.¹² In order to use this technique a pathophysiological approach that combines clinical, hemodynamic and echographic evaluation has been proposed¹³. It integrated oxygen transport/consumption, pressure-volume relationships and the concept of venous return.

Mechanism of hemodynamic instability

The various components of hemodynamic instability can be explained using the classical concept of venous return as described by Guyton.¹⁴ In simple terms, venous return (VR) is determined by a pressure gradient. This gradient corresponds to the difference between the mean systemic venous pressure (Pms) in the periphery and the right atrial pressure (Pra). This pressure gradient difference is divided by the resistance to venous return (Rvr).

$$VR = \frac{Pms - Pra}{Rvr} \quad (\text{Equation 1})$$

Therefore venous return and, consequently, cardiac output, will be reduced if: 1) the right atrial pressure is elevated, 2) the mean systemic pressure is low, and/or 3) the resistance to venous return is increased. There are several ways to illustrate this relationship. The classical approach to describe venous return and cardiac output is illustrated in Figure 1.¹⁵ The use of the pressure-volume relationship is used to describe a single cardiac cycle (Figure 2).




Figure 1 Venous return and cardiac output

The venous return and cardiac output (y axis) and its relation with right atrial pressure (x axis) is shown. The intersection of both curves will correspond to the right atrial pressure (Pra) at which, in a steady state, an individual will have an unique venous return and cardiac output. The mean systemic pressure (Pms) corresponds to the point where the venous return = 0. The venous return curve is linked to the resistance to venous return (Rvr) (dotted lines) (Adapted from Jacobsohn¹⁵).




Figure 2 Pressure and volume during a cardiac cycle
 (A) Changes in aortic, atrial, ventricular pressure, and ventricular volume in relation to the electrocardiogram. Left ventricular (LV) pressure and volume over time during a cardiac cycle is characterized by seven time-related events. Isovolumic contraction [1] is followed by early [2] and late [3] ejection. Diastole starts with isovolumic relaxation, [4] followed by the early filling phase after the opening of the mitral valve, [5] diastasis, [6] and atrial contraction. [7] (B) Corresponding LV pressure-volume relationship during one cardiac cycle (With permission of Denault *et al.* ¹⁶).

The pressure-volume relationship is typically described for the left ventricle but has also been used to evaluate right ventricular function. ¹⁷ The major difference between both ventricles is the reduced pressure in the right compared to the left ventricle. ¹⁸ In order to integrate the pressure-volume relationship to the venous return concept, we used a simplified alternative approach illustrated in figure 3. ¹⁹




Figure 3 Venous return and pressure-volume loop concept

The circulatory system can be divided into an intrathoracic and an extrathoracic compartment. Illustrated in red is the arterial system and in blue, the venous system. Most of the blood volume (~70%) lies in the venous system. The mean systemic pressure (Pms) is determined by the stressed volume (~30% of blood volume). The cardiac pump has two components, the right and the left ventricles, defined by their

respective pressure-volume loops (simplified). Both ventricles are connected together through the pericardium and the interventricular septum. The function and interaction between both ventricles will determine right atrial pressure (Pra). Blood returning back to the heart, or the venous return, will be dependent on the pressure gradient between the peripheral pressure, or Pms, and the central pressure, or Pra. Furthermore, any conditions increasing the resistance to venous return (Rvr), for instance compression of the inferior vena cava, will reduce venous return and consequently cardiac output. (Pa, systemic arterial pressure; Ra, arterial resistance)

The combination of conventional hemodynamic monitoring, FOCCUS \pm TEE allows the determination of the causes of hemodynamic instability.²⁰ However, so far, a systematic approach in the diagnosis of hemodynamic instability using conventional hemodynamic, FOCCUS \pm TEE could be useful in the OR, recovery room and ICU. This combined approach can be used to determine the causes of hemodynamic instability. The causes of hemodynamic instability resulting in reduced venous return or cardiac output and leading to difficult separation from CPB are a reduction in Pms, an increase in Pra and an increase in Rvr (Table 1).

Table 1 Mechanism of hemodynamic instability

1) Reduction in mean systemic pressure:

Reduction in stressed volume:

Hemorrhagic shock:

External hemorrhage

Internal: hemothorax, peritoneal hemorrhage, retroperitoneal hemorrhage, gastrointestinal hemorrhage

Increased in compliance

Drug-induced vasodilatation

Anaphylaxis

Vasoplegic syndrome

Sepsis

Adrenal insufficiency

2) Increased right atrial pressure

Left and right ventricular systolic dysfunction

Left and right ventricular diastolic dysfunction

Left and right outflow tract obstruction

Left and right embolism

Aortic and mitral patient-prosthesis mismatch

Hypoxia and hypercapnia

Pulmonary reperfusion syndrome

3) Increased resistance to venous return

Compartment syndrome

Pericardial tamponade

Mediastinal: post cardiopulmonary bypass

Pleural: hemothorax and pneumothorax

Abdominal: intrinsic, extrinsic or parietal

Vena cava syndrome

Inferior

Superior

Anesthesiologists are the critical care physicians of the operating and the recovery rooms. Echocardiography, both FOCCUS and TEE are becoming an integral part of the training curriculum of critical care physicians. Anesthesiologists are rapidly understanding the importance of acquiring these essential skills in resuscitation.

References

1. Rugolotto M, Chang CP, Hu B, Schnittger I, Liang DH. Clinical use of cardiac ultrasound performed with a hand-carried device in patients admitted for acute cardiac care. *Am J Cardiol* 2002;90:1040-2.
2. Gorcsan J, III, Pandey P, Sade LE. Influence of hand-carried ultrasound on bedside patient treatment decisions for consultative cardiology. *J Am Soc Echocardiogr* 2004;17:50-5.
3. Joseph MX, Disney PJ, Da Costa R, Hutchison SJ. Transthoracic echocardiography to identify or exclude cardiac cause of shock. *Chest* 2004;126:1592-7.
4. Manasia AR, Nagaraj HM, Kodali RB, Croft LB, Oropello JM, Kohli-Seth R, Leibowitz AB, DelGiudice R, Hufanda JF, Benjamin E, Goldman ME. Feasibility and potential clinical utility of goal-directed transthoracic echocardiography performed by noncardiologist intensivists using a small hand-carried device (SonoHeart) in critically ill patients. *J Cardiothorac Vasc Anesth* 2005;19:155-9.
5. Beaulieu Y, Marik PE. Bedside ultrasonography in the ICU: part 1. *Chest* 2005;128:881-95.
6. Beaulieu Y, Marik PE. Bedside ultrasonography in the ICU: part 2. *Chest* 2005;128:1766-81.
7. Beaulieu Y. Bedside echocardiography in the assessment of the critically ill. *Crit Care Med* 2007;35:S235-S249.
8. Lichtenstein DA, Meziere GA. Relevance of lung ultrasound in the diagnosis of acute respiratory failure: the BLUE protocol. *Chest* 2008;134:117-25.
9. Langlois SP. Focused ultrasound training for clinicians. *Crit Care Med* 2007;35:S138-S143.
10. Mazraeshahi RM, Farmer JC, Porembka DT. A suggested curriculum in echocardiography for critical care physicians. *Crit Care Med* 2007;35:S431-S433.
11. Price S, Via G, Sloth E, Guarracino F, Breikreutz R, Catena E, Talmor D. Echocardiography practice, training and accreditation in the intensive care: document for the World Interactive Network Focused on Critical Ultrasound (WINFOCUS). *Cardiovasc Ultrasound* 2008;6:49.
12. Mayo PH, Beaulieu Y, Doelken P, Feller-Kopman D, Harrod C, Kaplan A, Oropello J, Vieillard-Baron A, Axler O, Lichtenstein D, Maury E, Slama M, Vignon P. American College of Chest Physicians/La Société de Réanimation de Langue Française statement on competence in critical care ultrasonography. *Chest* 2009;135:1050-60.
13. Denault, A. Y. Difficult separation from cardiopulmonary bypass: importance, mechanism and prevention. 2010. Université de Montréal. Thesis/Dissertation (www.mediaenligne.umontreal.ca/medclin/anesth/denault_phd.wmv)
14. Guyton AC, Lindsey AW, Bernathy B, Richardson T. Venous return at various right atrial pressures and the normal venous return curve. *Am J Physiol* 1957;189:609-15.
15. Jacobsohn E, Chorn R, O'Connor M. The role of the vasculature in regulating venous return and cardiac output: historical and graphical approach. *Can J Anaesth* 1997;44:849-67.
16. Denault AY, Couture P, Tardif JC, Buithieu J. Transesophageal Echocardiography Multimedia Manual: A Perioperative Transdisciplinary Approach. Marcel Dekker, 2005.
17. Dell'Italia LJ, Walsh RA. Application of a time varying elastance model to right ventricular performance in man. *Cardiovasc Res* 1988;22:864-74.
18. Haddad F, Couture P, Tousignant C, Denault AY. The right ventricle in cardiac surgery, a perioperative perspective: I. Anatomy, physiology, and assessment. *Anesth Analg* 2009;108:407-21.
19. Deslauriers N, Dery R, Denault A. Acute abdominal compartment syndrome. *Can J Anaesth* 2009;56:678-82.
20. Denault AY, Couture P, Rochon A, Bouchard D. Échocardiographie transesophagienne au bloc opératoire. Conférences d'actualisation du Congrès national d'anesthésie et de réanimation 2007;1-16.